

NAXPET® Injectable solution.

ANALGESIC, ANTIPYRETIC AND NON-STEROIDAL ANTI-INFLAMMATORY.



Indicated for the treatment of feverish symptoms, inflammations and painful conditions of the bones, joints and musculoskeletal system in dogs and cats. Its use is especially recommended for the treatment of post-surgical pain, and in cases of osteoarthritis, inflammation of the musculoskeletal system and traumatic injuries in general.

Technical Specification

SPECIES

Dogs and cats.

DOSAGE FORM

Injectable solution.

THERAPEUTIC ACTION

Analgesic, Antipyretic and NON-STEROIDAL Anti-inflammatory.

COMPOSITION

Each mL of solution for injection contains:

Ketoprofen 10 mg

Excipients q.s.p 1 mL

INDICATIONS

Indicated for the treatment of feverish symptoms, inflammations and painful conditions of the bones, joints and musculoskeletal system in dogs and cats. Its use is especially recommended for the treatment of post-surgical pain, and in cases of osteoarthritis, inflammation of the musculoskeletal system and traumatic injuries in general.

ROUTE OF ADMINISTRATION AND DOSAGE

Route of administration:

- **Dogs:** Subcutaneous, intramuscular or intravenous route.
- **Cats:** Subcutaneous route.

Dose:

Dogs and cats:

0.2 mL per Kg of weight (2 mg / Kg), once a day, for up to 3 consecutive days.

CONTRAINDICATIONS

- Do not use in animals with known hypersensitivity to Ketoprofen.
- Do not use in animals with pathologies that present with gastrointestinal ulceration or bleeding.
- Do not use concomitantly with anticoagulants or corticosteroids.
- Do not use in animals with severe liver and kidney failure.
- Do not use concomitantly with other NSAIDs, or with diuretics.
- Do not use in pregnant females or in lactation period.

PRECAUTIONS

Avoid the use of the formulation in dehydrated, hypovolemic, hypotensive or hypoproteinemic animals since there is a greater potential risk of renal toxicity in them.

WARNINGS

Keep out of the reach of children.

ADVERSE EFFECTS

In some patients, edema, nausea, gastrointestinal irritation, headaches, nervousness, constipation, hives, skin rash or fever may be observed.

OBSERVATIONS

Special precautions for disposal of unused product or waste material:

- Discard any unused product remains in its original container.
- Dispose of the waste of this product with care together with household waste.

CONSERVATION

Store in a cool and dry place, at room temperature between 15 and 30 ° C.

CONDITION OF SALE

Sale with veterinary prescription.

PRESENTATION

20 mL vial

PREPARED BY

Laboratorio Drag Pharma Chile Invetec S.A.

RECORDS

Chile: Reg. SAG N° 0638

Uruguay: Reg. N° MGAP A-4441

Rep. Dominicana: Reg. N° 8862

Perú: Registro SENASA F.99.42.I.0075

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian