
# **NAXPET<sup>®</sup> Oral suspension**

#### ANALGESIC, ANTIPYRETIC AND NON-STEROIDAL ANTI-INFLAMMATORY



Indicated for the treatment of feverish pictures, inflammations and painful conditions of the bones, joints and skeletal muscle system in dogs and cats. Its use is especially recommended for the treatment of post-surgical pain, and in cases of osteoarthritis, inflammation of the musculoskeletal system and traumatic injuries in general.

# **Technical Specification**

**SPECIES** 

Dogs and cats.

**DOSAGE FORM** 

Oral suspension

THERAPEUTIC ACTION

Analgesic, Antipyretic and Non-steroidal anti-inflammatory

**COMPOSITION** 

# **INDICATIONS**

Indicated for the treatment of feverish pictures, inflammations and painful conditions of the bones, joints and skeletal muscle system in dogs and cats. Its use is especially recommended for the treatment of post-surgical pain, and in cases of osteoarthritis, inflammation of the musculoskeletal system and traumatic injuries in general.

#### ROUTE OD ADMINISTRATION AND DOSAGE

Administration route: Oral administration.

#### Dose

1 mg / Kg once a day for 5 days, (equivalent to 1 mL for every 4 kilos of weight or 8 drops per Kg of weight, once a day for 5 days).

In the case of post-operative pain and chronic pain, administer an initial dose of 2 mg / Kg and then 1 mg / Kg once a day for 5 days (equivalent to an initial dose of 2 mL every 4 Kg of weight or 16 drops per Kg of weight and then 1 mL every 4 Kg of weight or 8 drops per Kg of weight, once a day for 5 days).

#### **CONTRAINDICATIONS**

- Do not use in animals with known hypersensitivity to Ketoprofen.
- Do not use in animals with pathologies that present with ulceration or gastrointestinal bleeding.
- Do not use in animals with severe kidney and liver failure.
- Do not use concomitantly with anticoagulants or corticosteroids.
- Do not use concomitantly with other NSAIDs.
- Do not use in pregnant or lactating females.

#### SPECIAL PRECAUTIONS FOR USE

# Warning and special precautions for use:

- To avoid the risk of increased adverse effects in dogs and cats, it is recommended not to administer the formulation in conjunction with other non-steroidal anti-inflammatory drugs.
- Ketoprofen should be used with caution in animals that present gastrointestinal pathologies or chronic inflammatory bowel disease, and should not be used if there is gastric ulceration or gastrointestinal bleeding.
- Avoid using the formulation in dehydrated, hypovolemic, hypotensive or hypoproteinemic animals, since there is a greater potential risk of kidney toxicity in them.
- Keep out of the reach of children.

# SPECIAL PRECAUTIONS FOR THE OPERATOR

- After handling and applying the product, a hand wash is recommended.
- Avoid contact of the product with the skin and with the eyes.
- In case of accidental ingestion of the product, it is recommended to go to a medical center and show the internal brochure of the product.

#### ADVERSE EFFECTS

Ketoprofen can cause gastrointestinal disturbances at therapeutic doses in some animals. It can cause vomit, diarrhea and transient anorexia in dogs and cats.

# **OBSERVATIONS**

# Special precautions for the disposal of unused product or waste material:

Empty containers can be discarded as household waste, without any special precautions. Do not dispose of containers with product residues on the ground or water courses. For expired or unused products, contact the manufacturing laboratory.

#### **CONSERVATION**

- Store in a dry place protected from light, at room temperature between 15 and 30 °C.
- Once opened, use the product within 3 months. Discard the unused product after that period of time.

# **CONDITION OF SALE**

Sale with a Veterinary Medical prescription only .

#### **PRESENTATION**

# 20 mL bottle

## PREPARED BY

Laboratorio Drag Pharma Chile Invetec S.A.

# **RECORDS**

Chile: Reg. SAG N° 1995

Costa Rica: Reg. N° MAG CL 4-14-4-6177 Uruguay: Reg. MGAP Nº 2018A00608 Perú: Registro SENASA F.C6.02.I.0001

## **COUNTRIES WHERE IT IS MARKETED**

Importer in Uruguay:

VIVAFIL S.A.

Rio Negro 1107 Montevideo - Uruguay, TEL 29001112

grupotecnovet@gmail.com

Technical Director: DMTV Diego Cuadrado.

Imported and Distributed in Peru by:

Representaciones Durand SAC.

Av. Manuel Olguín N ° 501 Office N ° 604 Santiago de Surco Lima.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian