

REGEPIPEL® PLUS - Shampoo

ANTIFUNGAL AND ANTISEPTIC SHAMPOO.



For the treatment of seborrheic dermatitis associated with *Staphylococcus intermedius* infections, *Malassezia pachydermatis*, and as an aid in the treatment of dermatophyte infections due to *Microsporum canis* in dogs and cats.

Technical Specification

SPECIES

Dogs and cats.

DOSAGE FORM

Shampoo.

THERAPEUTIC ACTION

Antifungal and antiseptic shampoo.

COMPOSITION

Each 100 mL contains:
Chlorhexidine Gluconate 20% 10 g
(Equivalent to 2 g of Chlorhexidine Gluconate)
Miconazole Nitrate 2.3 g
(Equivalent to 2 g of Miconazole base)
Excipients q.s.p 100 mL

INDICATIONS

For the treatment of seborrheic dermatitis associated with *Staphylococcus intermedius* infections, *Malassezia pachydermatis*, and as an aid in the treatment of dermatophyte infections due to *Microsporum canis* in dogs and cats.

MODE OF APPLICATION

- Wet the animal with plenty of water.
- Apply a sufficient amount of the shampoo at various points and massage. With the help of a

sponge, spread to promote its penetration into the skin and coat. Be sure to apply Regepipel® Plus around the lips, under the tail and between the fingers as these are reservoir areas for causative agents.

- Leave on for 10 to 15 minutes, then rinse with plenty of clean water.
- Repeat two or three times a week until symptoms subside and then once a week or as directed by the Veterinary Doctor.

ROUTE OF ADMINISTRATION AND DOSAGE

Topically.

DRUG INTERACTIONS

Chlorhexidine is inactivated when used with common detergents or anionic compounds.

CONTRAINDICATIONS

- Do not use in cats younger than 6 weeks of age and in dogs younger than 4 weeks of age.
- Do not administer to individuals with known hypersensitivity to any component of the preparation.
- Do not use in pregnant or lactating females.

PRECAUTIONS

Use in cats older than 6 weeks old and in dogs older than 4 weeks old.

SPECIAL PRECAUTIONS FOR THE OPERATOR

Avoid contact with eyes, in case of accidental contact wash immediately with plenty of water.

WARNINGS

Special warnings and precautions for use:

- In the treatment of dermatophytosis, the association of topical therapy with systemic therapy favors the healing of the lesions in 6-9 weeks, less time than required when only systemic therapy is administered.
- In case of local irritation, stop treatment.
- Avoid contact with the eyes of the animal, in case of accidental contact wash immediately with plenty of water.
- Keep out of the reach of children.

OBSERVATIONS

Special precautions for the disposal of unused product or waste material:

Empty containers can be discarded as household waste, without any special precautions. Do not dispose of containers with product residues on the ground or water courses. For expired or unused products, contact the manufacturing laboratory.

CONSERVATION

Keep at room temperature, between 15 and 30°C.

CONDITION OF SALE

To be supplied only on veterinary prescription.

PRESENTATION

Bottle with 150 mL

PREPARED BY

Laboratorio Drag Pharma Chile Invetec S.A.

RECORDS

Chile: Reg. SAG N° 1696

Panamá: Reg. N° RF-8355-18

El Salvador: VE2013084783

Bolivia: Reg. SENASAG PUV-F N° 007259/16

Uruguay: Reg. MGAP N° 2019A00682

COUNTRIES WHERE IT IS MARKETED

Distribution in El Salvador: Rafael Alfredo Alfaro Castillo.
8a C. Pte. Pje. Moreno N ° 112, Col. Flor Blanca.
San Salvador, El Salvador.

Imported and distributed in Bolivia by:

ZOOFARMA

TEL: +(591)222-3357

Street Díaz Romero 1339, La Paz.

Imported in Uruguay by: VIVAFIL S.A.

RIO NEGRO 1107 Montevideo - Uruguay, TEL 29001112

grupotecnovet@gmail.com

Technical Director: DMTV Diego Cuadrado.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian