# **RESPIG®** Oral suspension

#### **BROAD SPECTRUM ANTIMICROBIAL**



Broad spectrum antimicrobial indicated for the treatment of respiratory tract and urinary tract infections and soft tissue infections caused by Gram positive and Gram negative bacteria susceptible to Sulfamethoxazole and Trimethoprim.

## **Technical Specification**

**SPECIES** 

Dogs and cats.

**DOSAGE FORM** 

Oral suspension

THERAPEUTIC ACTION

Broad spectrum antimicrobial

## **COMPOSITION**

## **INDICATIONS**

Broad spectrum antimicrobial indicated for the treatment of respiratory tract and urinary tract infections and soft tissue infections caused by Gram positive and Gram negative bacteria susceptible to Sulfamethoxazole and Trimethoprim.

#### ROUTE OD ADMINISTRATION AND DOSAGE

## Orally.

## • Dosage of active ingredients:

30 mg of the association per kilo of weight every 24 hours, or 15 mg of the association per kilo of weight every 12 hours, for 7 to 10 days.

• Product dosage:

5 mL for every 8 Kg of weight, every 24 hours; or 2.5 mL every 8 Kg of weight, every 12 hours, for 7 to 10 days.

Do not administer for more than 14 days.

## **DRUG INTERACTIONS**

- The association of Sulfamethoxazole and Trimethoprim may prolong clotting times in patients medicated with coumarin anticoagulants.
- Antacids can decrease the bioavailability of sulfonamides if administered concurrently.
- Trimethoprim can reduce the therapeutic effects of cyclosporine (systemic) and increase the risk of nephrotoxicity.

#### **CONTRAINDICATIONS**

- Do not use in dogs with liver and / or kidney damage.
- Do not use in dogs with hypersensitivity to sulfonamides.
- Do not use in Pinscher breed dogs.
- Do not use in pregnant or lactating females.

#### SPECIAL PRECAUTIONS FOR THE OPERATOR

Do not handle by people hypersensitive to sulfonamides.

#### **WARNINGS**

Keep out of the reach of children.

#### **OBSERVATIONS**

## Special precautions for disposal of unused product or waste material:

Empty containers can be disposed of as household waste, without any special precautions. Do not dispose of containers with product remains on the ground or water courses. For expired or unused products, contact the manufacturing laboratory.

#### **CONSERVATION**

Store between 15 and 30 °C, protected from light.

Once the container is opened, use within 14 days. Discard the unused product after that period of time

#### **CONDITION OF SALE**

Sale with veterinary prescription.

#### **PRESENTATION**

100 mL bottle

## PREPARED BY

Laboratorio Drag Pharma Chile Invetec S.A.

#### **RECORDS**

Chile: Reg. SAG N° 944

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian