

RUMITEN® Oral suspension

BROAD SPECTRUM ANTIPARASITIC



Indicated for the treatment of gastrointestinal and pulmonary parasitosis in sheep and goats caused by parasites sensitive to the combination of Fenbendazole and Praziquantel, such as: *Moniezia spp.*, Mature and immature forms of *Haemonchus spp.*, *Cooperia spp.*, *Nematodirus spp.*, *Chabertia ovina*, *Oesophagostomum spp.*, *Trichostrongylus spp.*, *Trichuris ovis.*, *Ostertagia ostertagi* (hypobiotic adults and larvae), *Ostertagia circumcincta* and *Dictyocaulus filaria*. It is recommended to treat all the animals in the herd every three months. This contributes to reducing the pressure of parasitic infection, achieving epidemiological control.

Technical Specification

SPECIES

Sheep and goats.

DOSAGE FORM

Oral suspension

THERAPEUTIC ACTION

Broad spectrum antiparasitic

COMPOSITION

Each 100 mL of oral suspension contains:

Fenbendazole 2 g
Praziquantel 1.5 g
Excipients q.s.p 100 mL

INDICATIONS

Indicated for the treatment of gastrointestinal and pulmonary parasitosis in sheep and goats caused by parasites sensitive to the combination of Fenbendazole and Praziquantel, such as: *Moniezia spp.*, Mature and immature forms of *Haemonchus spp.*, *Cooperia spp.*, *Nematodirus spp.*, *Chabertia ovina*, *Oesophagostomum spp.*, *Trichostrongylus spp.*, *Trichuris ovis.*, *Ostertagia ostertagi* (hypobiotic adults and larvae), *Ostertagia circumcincta* and *Dictyocaulus filaria*. It is recommended to treat all the animals in the herd every three months. This contributes to reducing the pressure of parasitic infection, achieving epidemiological control.

EFFECTIVENESS

Efficacy period: 36 months.

ROUTE OF ADMINISTRATION AND DOSAGE

Oral administration.

Recommended practice dose:

2.5 mL per 10 kg of body weight in a single dose (equivalent to a dose of 3.75 mg of Praziquantel and 5 mg of Fenbendazole per kg of live weight).

CONTRAINDICATIONS

Do not administer to animals with known hypersensitivity to any active ingredient.

WARNINGS

- Shake before using.
- Keep out of the reach of children.
- Only for oral use in sheep and goats.

GUARD PERIOD

Meat: 14 days.

Milk: 7 days.

OBSERVATIONS

Special precautions for disposal of unused product or waste material:

Discard any unused product remains in its original container. Do not throw the empty container or with product remains, in rivers, lakes or streams of natural water. Do not reuse the container. Dispose of this product waste carefully with household waste.

CONSERVATION

Store at a temperature between 2 and 30 ° C, protected from light.
Once the container is opened, use within 1 month. Discard the unused product after that period.

CONDITION OF SALE

To be supplied only on veterinary prescription.

PRESENTATION

Bottle with 100 mL, 1 Liter and 3 Liters.

PREPARED BY

Laboratorio Drag Pharma Chile Invetec S.A.

RECORDS

Reg. SAG Nº 1961

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian