

SINPUL[®] SPRAY Solution

EXTERNAL ANTIPARASITICS.



Eliminates fleas (*Ctenocephalides felis*) and ticks (*Rhipicephalus sanguineus*) in dogs.

Technical Specification

SPECIES

Dogs.

DOSAGE FORM

Spray solution.

THERAPEUTIC ACTION

External Antiparasitics.

COMPOSITION

Each 100 mL contains:

Permethrin.....0.2 g
Piperonyl Butoxide.....2.0 g
Excipients q.s.....100.0 mL

INDICATIONS

Eliminates fleas (*Ctenocephalides felis*) and ticks (*Rhipicephalus sanguineus*) in dogs.

MODE OF APPLICATION

Spray the product all over the dog's body, from the tail to the neck (application against the grain). After applying, rub the entire body vigorously with your hands so that the product achieves maximum contact with the skin and not only remains on the coat. If the dog has a large number of fleas and / or ticks, it is recommended to repeat the application every 7 to 15 days.

ROUTE OF ADMINISTRATION AND DOSAGE

Topical use.

Dosage of active ingredients:

6 to 18 mg / Kg of Permethrin and 60 to 80 mg / Kg of Piperonyl Butoxide.

Product dosage:

Apply 4 to 11 sprays per kg of weight, depending on the length of the coat, in a single dose.

CONTRAINDICATIONS

- Do not use in cats.
- Do not use in puppies under 6 weeks of age.
- Do not administer in animals with hypersensitivity to the active principles.
- Do not administer in pregnant or lactating females.

SPECIAL PRECAUTIONS FOR USE

- Flammable product.
- Avoid contact with the eyes, mouth or other mucous membranes.
- In case of irritation, stop using the product and wash with plenty of water.

SPECIAL PRECAUTIONS FOR THE OPERATOR

- When applying the product wear protective gloves.
- Wash your hands with plenty of water after use.
- In case of contact with the eyes, mouth or skin, wash the affected area with plenty of water.

WARNINGS

- Keep out of the reach of children.
- External use.

ADVERSE EFFECTS

Hypersensitivity reactions have been described when Permethrin is administered topically. In some cases, itching or mild irritation may occur at the application site.

OBSERVATIONS

Special precautions for disposal of unused product or waste material:

Discard the remains of unused product in its original container, well closed. Do not throw the empty container or with product remains, in soils, rivers, lakes or streams of water, as it is toxic to aquatic animals. Do not reuse the container. Dispose of this product waste carefully with household waste.

CONSERVATION

Keep at room temperature (between 15 and 30 ° C), in a dry place and away from fire. Once opened, use the product within 12 weeks. Discard the unused product after that period of time.

CONDITION OF SALE

OTC product (non-prescription).

PRESENTATION

200 mL and 500 mL spray bottle

PREPARED BY

Drag Pharma Laboratory.

RECORDS

Reg. SAG N° 0390

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian