EQUIMIC® PLUS - Oral Paste

BROAD SPECTRUM ANTI-PARASITE.



Equimic[®] Plus is a broad spectrum antiparasitic. because of combination of Ivermectin and Praziquantel is effective against large strongyles (Strongylus spp, Triodontophorus spp.), small strongyles (Cylicocyclus spp, Cylicostephanusspp, Cylicodontophorus spp, Gyalocephalus spp.), pinworms (Oxyuris equi), ascarids (Parascaris equorum), trichostrongyles (*Trichostrongylus axei*), intestinal nematodes (Strongyloides westeri), stomach nematodes (Habronema spp., Draschia megastoma), horse bot fly (Gasterophilus spp.) and tapeworms (Anoplocephala perfoliata).

Technical Specification

SPECIES

Horses.

DOSAGE FORM

Oral paste.

THERAPEUTIC ACTION

Broad spectrum anti-parasite.

COMPOSITION

INDICATIONS

Equimic[®] Plus is a broad spectrum antiparasitic, because of the combination of Ivermectin and Praziquantel is effective against large strongyles (*Strongylus spp, Triodontophorus spp.*), small strongyles (*Cylicocyclus spp, Cylicostephanus spp, Cylicodontophorus spp, Gyalocephalus spp.*), pinworms (*Oxyuris equi*), ascarids (*Parascaris equorum*), trichostrongyles (*Trichostrongylus axei*), intestinal nematodes (*Strongyloides westeri*), stomach nematodes (*Habronema spp., Draschia megastoma*), horse bot fly (*Gasterophilus spp.*) and tapeworms (*Anoplocephala perfoliata*).

Active ingredient dose:

• Ivermectin: 0.2 mg/Kg, single dose.

• Praziquantel: 1.5 mg/Kg, single dose.

Product dose:

3.3 g of Equimic[®] Plus for each 150 Kg of body weight, single dose. Each syringe is enough for up to 900 kg of weight.

Place the dosing Equimic[®] Plus syringe directly into the horse's mouth and download the volume equivalent to the recommended dosage according to the weight of the animal.

CONTRAINDICATIONS

- Do not use in case of hypersensitivity to any of the active ingredients.
- Do not use in foals under 2 month old and stallions.

SPECIAL PRECAUTIONS FOR USE

Equimic[®] Plus has been formulated to be used exclusively in horses. It should not be used in other species.

SPECIAL PRECAUTIONS FOR THE OPERATOR

- In the event of accidental ingestion, do not induce vomiting. Consult to the doctor immediately.
- If contact with skin or eyes, wash with abundant water. If irritation occurs, consult to the doctor.
- Wash your hands after the administration of this product.

WARNINGS

Keep out of reach of children.

ADVERSE EFFECTS

In the case of serious infestations, it can be observed signs of transient colic.

GUARD PERIOD

Do not administer to horses which meat is intended for human consumption.

OBSERVATIONS

Special precautions for disposal of unused product or waste material:

Do not dispose product container, empty or with product along with the domestic waste. Do not eliminate it on the ground or in water courses since the product is dangerous to aquatic organisms.

CONSERVATION

Store in a cool and dry place, between 15° and 30°C, protected from sunlight. Once the syringe has been opened, use within 3 months.

CONDITION OF SALE

To be supply only on veterinary prescription.

PRESENTATION

20 g graduated syringe.

RECORDS

Chile: Reg. SAG No. 2171-B

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian