

# INVECLOR® - Antiseptic soap

---

ANTISEPTIC SOAP FOR SKIN AND MUCOUS MEMBRANES.



INVECLOR®, Chlorhexidine Gluconate 2%, liquid soap, is indicated as an antiseptic soap for skin and mucous membranes.

## Technical Specification

### SPECIES

---

Horses

### DESCRIPTION

---

The INVECLOR® product combines the advantages of soapy cleaners with an effective, broad-spectrum and safe antiseptic such as Chlorhexidine, making it ideal for treating skin and / or mucous areas where it is required to reduce contamination quickly and persistent.

Chlorhexidine has a broad spectrum of antimicrobial activity. It is very effective against Gram positive bacteria, especially *Staphylococcus aureus*, other staphylococci, and various species of streptococci. It is less effective against Gram negatives such as *Escherichia coli*, *Klebsiella*, *Salmonella*, and *Pseudomonas*. It is capable of preventing the germination of bacterial spores. It is moderately active against *M. tuberculosis*. It is active against fungi such as *Candida albicans*, *Microsporum gypseum*, *M. canis* and *Trichophyton mentagrophytes* and yeasts. It has variable activity against viruses.

### DOSAGE FORM

---

Liquid soap. Chlorhexidine Gluconate 2%.

### THERAPEUTIC ACTION

---

Antiseptic soap for skin and mucous membranes.

### COMPOSITION

---

Each 100 mL of soap contains:  
Chlorhexidine Gluconate ..... 2 g  
Excipients q.s.p ..... 100 mL

### INDICATIONS

---

INVECLOR®, Chlorhexidine Gluconate 2%, liquid soap, is indicated as an antiseptic soap for skin and mucous membranes.

#### MODE OF APPLICATION

---

Before applying this product, remove dirt and contamination from the area to be treated. Then, moisten the area and apply the product directly on the skin and / or mucous membranes in sufficient quantity so that the area is completely covered. Scrub (with a brush, brush, sponge or by hand) the area gently for 5 minutes.

Rinse with plenty of water to remove the entire product and dry with a towel, gauze or sterile compress.

#### ROUTE OF ADMINISTRATION AND DOSAGE

---

Topical use.

Place a large quantity of INVECLOR®, Chlorhexidine Gluconate 2%, liquid soap, on the skin and / or mucosa area to be treated so that it is completely covered by the product.

#### DRUG INTERACTIONS

---

Do not mix with other disinfectants and antiseptics.

#### CONTRAINDICATIONS

---

- Do not use in animals that are hypersensitive to chlorhexidine gluconate.
- Contraindicated in cases of animals with wounds, since its use delays the healing process.

#### SPECIAL PRECAUTIONS FOR USE

---

- The applied product should be allowed to act for 5 minutes before rinsing.
- To improve antiseptic conditions, it is recommended to depilate the treated area before using this product.
- Do not use in the eyes, ears, or inside the animal's mouth; In case of contact wash with a lot of water.

#### SPECIAL PRECAUTIONS FOR THE OPERATOR

---

- It should not be handled by people who are hypersensitive to Chlorhexidine.
- Wash hands after use. The use of gloves is recommended.
- In case of contact with the skin or the eyes, it is recommended to wash with plenty of water. If irritation develops and persists, see doctor.

#### WARNINGS

---

Keep out of the reach of children.

#### OBSERVATIONS

---

##### **Use during pregnancy and lactation:**

Indicated for safe use in pregnant or lactating females.

#### CONSERVATION

---

Store in a dry place at temperatures between 2 and 30° C protected from light.

#### CONDITION OF SALE

---

Free sale.

VETERINARY USE.

EXTERNAL USE.

PRESENTATION

---

1 liter bottle

PREPARED BY

---

Laboratorio Drag Pharma.

RECORDS

---

Reg. SAG Nº 2242.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian