

BILIFAR® Oral powder

RUMINAL DIGESTION STIMULANT



BILIFAR® Oral Powder, is a stimulant of ruminal digestion. Its use is recommended in pathologies in which it is necessary to stimulate ruminal fermentation and / or increase blood glucose levels, such as simple ruminal indigestion, alkalosis or ruminal rot, ketosis adjuvant and / or sudden changes in diet. BILIFAR® is also an alternative source of minerals for ruminants.

Technical Specification

SPECIES

Cattle, sheep and goats.

DOSAGE FORM

Oral powder

THERAPEUTIC ACTION

Ruminal Digestion Stimulant

COMPOSITION

Each bag with 120 grams contains:

Sodium propionate	60 g
Calcium propionate	40 g
Sodium Chloride	18.5 g
Cobalt Chloride	40 mg
Copper Sulfate	150 mg
Manganese Sulfate	200 mg
Iron Sulfate	300 mg
Zinc Chloride	10 mg
Excipients q.s.p	120 grams

INDICATIONS

BILIFAR® Oral Powder, is a stimulant of ruminal digestion. Its use is recommended in pathologies in which it is necessary to stimulate ruminal fermentation and / or increase blood glucose levels, such as simple ruminal indigestion, alkalosis or ruminal rot, ketosis adjuvant and / or sudden changes in diet. BILIFAR® is also an alternative source of minerals for ruminants.

ROUTE OF ADMINISTRATION AND DOSAGE

Administration route:
Oral.

Product dosage:

- Adult cattle (cows, oxen, bulls) 450 to 600 Kg: 1 to 2 bags of 120 g dissolved in 1 to 3 liters of water.
- Young cattle (calves, steers) 150 to 300 Kg: 1/2 bag of 120 g dissolved in 1/2 to 1 liter of Water.
- Sheep and goats: 1/4 bag of 120 g dissolved in 1/4 liter of water.

The treatment can be repeated every 12 to 24 hours, for up to 10 days.

CONTRAINDICATIONS

- Do not use in case of hypersensitivity to some components of the formula.
- Do not administer to dehydrated animals without prior hydration therapy.

WARNINGS

- Keep out of the reach of children.
- Respect the indicated dose considering the safety margin of Copper Sulfate.

GUARD PERIOD

Meat: 0 days.
Milk: 0 days.

OBSERVATIONS

Overdose:

Symptoms of Copper toxicity include hemolysis, hemoglobinuria, and jaundice. The oral toxic dose of Copper for sheep and calves is 20 to 110 mg / Kg; for cattle (adults) it is from 220 to 880 mg / Kg; and for goats it is 60 mg / Kg.

Special precautions for the disposal of unused product or waste material:

Eliminate the remains of unused product in its original container. Do not throw the empty container or with product remains, in rivers, lakes or torrents of water. Dispose of this product with caution with household waste.

CONSERVATION

Store in a cool, dry place, at room temperature between 15° and 30°C. Use immediately once prepared and discard the excess product.

CONDITION OF SALE

Sale only with a Veterinary Medical prescription.

PRESENTATION

Sachet with 120 grams.

RECORDS

- Chile: Reg. SA. N° 280

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian