BOVIFORT[®] Injectable solution

BROAD SPECTRUM ANTIPARASITIC - FASCIOLICIDA



Bovifort[®] is a broad-spectrum injectable antiparasitic that is indicated for the treatment of infestations by gastrointestinal, pulmonary and ectoparasites in cattle. Due to its specific action as a fasciolicide, it is also indicated in the treatment of hepatic dystomatosis, caused by adult Fasciola hepatica, also exercising adequate control over juvenile stages of 8 weeks or more. Its use is indicated in the treatment and control of the main internal and external parasites of bovines such as: Ostertagia sp .; Haemonchus sp .; Trichostrongylus sp .; Cooperia sp .; Oesophagostomum radiatum; Nematodirus sp.; Dictyocaulus viviparus; Fasciola hepatica; Haematopinus eurysternus; Linognathus vituli ,; Psoroptes bovis, Sarcoptes scabiei var bovis.

Technical Specification

SPECIES

Bovines

DOSAGE FORM

Injectable solution

THERAPEUTIC ACTION

Broad spectrum antiparasitic - Fasciolicida

COMPOSITION

Each 1 mL of solution for injection contains: Ivermectin 10 mg Clorsulon 100 mg Excipients q.s.p 1 mL

INDICATIONS

Bovifort[®] is a broad-spectrum injectable antiparasitic that is indicated for the treatment of infestations by gastrointestinal, pulmonary and ectoparasites in cattle. Due to its specific action as a fasciolicide, it is also indicated in the treatment of hepatic dystomatosis, caused by adult Fasciola

hepatica, also exercising adequate control over juvenile stages of 8 weeks or more. Its use is indicated in the treatment and control of the main internal and external parasites of bovines such as: Ostertagia sp .; Haemonchus sp .; Trichostrongylus sp .; Cooperia sp .; Oesophagostomum radiatum; Nematodirus sp .; Dictyocaulus viviparus; Fasciola hepatica; Haematopinus eurysternus; Linognathus vituli ,; Psoroptes bovis, Sarcoptes scabiei var bovis.

ROUTE OD ADMINISTRATION AND DOSAGE

Route of administration:

Subcutaneous. Do not apply the product intravenously or intramuscularly. In some cases there may be mild pain at the injection site.

Dosage of active ingredients: Ivermectin: 0.2 mg / Kg and Chlorsulon: 2 mg / Kg Single dose.

Product dosage: 1 mL for every 50 Kg of weight in a single dose.

WARNINGS

Keep out of the reach of children

GUARD PERIOD

Beef cattle: 49 days.

Milk cattle: Do not administer to animals whose milk is intended for human consumption.

CONSERVATION

Keep at room temperature between 15 ° and 30 ° C, in a dry place protected from light.

CONDITION OF SALE

Sale with veterinary prescription.

PRESENTATION

50 mL ampoule bottle and 500 mL serum bottle

PREPARED BY

Laboratorio Drag Pharma Chile Invetec S.A.

RECORDS

Chile: Reg. SAG N° 1880-B

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian