# **DIARREPAS**<sup>®</sup> Oral suspension

## ANTIDIARRHEAL AND CHEMOTHERAPY.



It is recommended for use in intestinal infections caused bν sensitive non-absorbable to sulfonamides organisms. recommended for the treatment of enteric infections caused by gram positive and aram negative organisms. It has therapeutic use in acute and chronic diarrhea cases in dogs and cats, since it has protective mucosal chemotherapeutic action that avoids injury in the intestinal epithelium caused by irritating substances. By the presence of intestinal adsorbents, also prevent the absorption of toxins, gases or other toxic substances of organic or inorganic origins that can complicate a diarrheal case.

## **Technical Specification**

#### **SPECIES**

Dogs and cats.

**DOSAGE FORM** 

Oral suspension.

THERAPEUTIC ACTION

Antidiarrheal and chemotherapy.

## **COMPOSITION**

## **INDICATIONS**

It is recommended for use in intestinal infections caused by sensitive to non-absorbable sulfonamides organisms. It is recommended for the treatment of enteric infections caused by gram positive and gram negative organisms. It has therapeutic use in acute and chronic diarrhea cases in dogs and cats, since it has protective mucosal chemotherapeutic action that avoids injury in the intestinal epithelium caused by irritating substances. By the presence of intestinal adsorbents, also prevent the absorption of toxins, gases or other toxic substances of organic or inorganic origins that

can complicate a diarrheal case.

## **ROUTE OD ADMINISTRATION AND DOSAGE**

## Oral route.

- **Dosage of the active ingredients:** Phthalsulfatiazole 100 mg / Kg; Sulfaguanidine 100 mg / Kg; Colloidal Kaolin 200 mg / Kg, Citric Pectin 10 mg / Kg, every 12 hours for 4 days.
- Product dose: 10 mL for every 10 kilos of weight, every 12 hours for 4 days.

### **CONTRAINDICATIONS**

- Do not use simultaneously or associated with other chemotherapeutic agents, or other substances with similar action.
- Do not administer to animals showing resistance to the sulfonamides treatment.
- It should not be used in animals presenting a history of hypersensitivity to some component of the product.

#### **WARNINGS**

Keep out of reach of children.

#### **OBSERVATIONS**

Shake before using.

## Special precautions for the disposal of unused product or waste material:

Discard the remains of unused product in its original container. Do not throw the empty container or with product remains, in rivers, lakes or streams of natural water. Dispose of this product with caution with household waste.

## **CONSERVATION**

Store at a temperature between 15 and 30 ° C, protected from light. Once opened, use the product within 7 days. Discard the unused product after that period of time.

## **CONDITION OF SALE**

To be supply only on veterinary prescription.

## **PRESENTATION**

Bottle containing 100 mL

#### PREPARED BY

Drag Pharma Laboratory.

#### **RECORDS**

Reg. SAG No. 133

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of

this information without consulting a veterinarian