

# FIPRODRAG® 0,5 mL - Cats

## TOPICAL SOLUTION.



FIPRODRAG® is indicated for the treatment and control of fleas (*Ctenocephalides felis*) and mites (*Cheyletiella spp.*) infestation in cats. FIPRODRAG® is also recommended as adjuvant therapy in flea allergy dermatitis (FAD) in cats.

**Efficacy:** With a single application, FIPRODRAG® allows to control flea reinfestation in cats, during a period of at least 5 weeks. If parasite burden is high, a monthly application of product is advisable.

For its lipophilic properties, FIPRODRAG® resists baths carried out 48 hours after its application. Do not bathe your pet 48 hours before and after applying the product.

## Technical Specification

### DOSAGE FORM

Topical solution.

### THERAPEUTIC ACTION

Topical solution.

### COMPOSITION

Each mL of solution contains:

Fipronil.....100 mg

Excipients q.s.p.....1 mL

### PROPERTIES

Effective action against fleas in cats.

### INDICATIONS

FIPRODRAG® is indicated for the treatment and control of fleas (*Ctenocephalides felis*) and mites (*Cheyletiella spp.*) infestation in cats. FIPRODRAG® is also recommended as adjuvant therapy in flea allergy dermatitis (FAD) in cats.

**Efficacy:** With a single application, FIPRODRAG® allows to control flea reinfestation in cats, during a period of at least 5 weeks. If parasite burden is high, a monthly application of product is advisable.

For its lipophilic properties, FIPRODRAG® resists baths carried out 48 hours after its application. Do not bathe your pet 48 hours before and after applying the product.

#### MODE OF APPLICATION

---

1. Remove the upper marked tip of the pipette.
2. Put the end of the pipette directly on the animal skin, at scapulae (shoulders) level or at the base of neck, trying to separate the hair.
3. Press and pour the full content of pipette on 2 or 3 points, as indicated in the figure.

#### ROUTE OF ADMINISTRATION AND DOSAGE

---

For external use, through skin application.

Dose of active ingredient:

From 6.7 to 13.4 mg/kg body weight of the animal to be treated. In cats, apply one pipette of 0.5 mL.

#### CONTRAINDICATIONS

---

- Do not use on kittens less than 12 weeks old.
- Do not use in rabbits.

#### PRECAUTIONS

---

Administer with caution in weakened animals.

#### SPECIAL PRECAUTIONS FOR THE OPERATOR

---

##### **Precautions to be taken by the person administering the product:**

- Avoid touching the pipette content directly with your fingers.
- Wash your hands with soap and water after applying and handling the product.
- Do not eat food, smoke, or drink beverages during the application of product.
- Avoid handling your pet immediately after the treatment, until the coat is completely dry.
- Discard in a safe way the empty pipettes. Do not dispose the empty pipettes into natural waterways.

#### WARNINGS

---

- Keep out of reach of children.
- For external use on cats only.
- The product may be harmful if eaten.
- The product is toxic to fishes, birds and bees.
- Flammable product.

#### SIDE EFFECTS

---

If the animal licks itself, a short period of hypersalivation may occur. If there is hypersensitivity or local irritation, it is recommended to not repeat the treatment.

#### OBSERVATIONS

---

- POISON.
- Flammable.

#### CONSERVATION

---

- Store in the original pack, at room temperature between 15 and 30°C, and in a fresh place.
- Keep out of heat sources.
- Do not store with food products.

#### CONDITION OF SALE

---

OTC product (non-prescription).

#### PRESENTATION

---

Box or Thermo-contracting Display containing 1 pipette of 0.5 mL

#### PREPARED BY

---

Drag Pharma Laboratory.

#### RECORDS

---

SAG Reg. No. 2162

#### COUNTRIES WHERE IT IS MARKETED

---

Imported and distributed in Bolivia by:

ZOOFARMA  
TEL: +(591)222-3357  
Street Díaz Romero 1339, La Paz.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian