

FIPROKILL[®] SPRAY - Spray Solution

EXTERNAL ANTIPARASITIC.



FIPROKILL[®] SPRAY is effective in the control and treatment of flea (*Ctenocephalides felis*) and tick (*Rhipicephalus sanguineus*) infestation in dogs and cats from 2 days of age, providing a rapid knock-down effect against adult fleas.

Technical Specification

SPECIES

Dogs and cats.

DOSAGE FORM

Topical Solution

THERAPEUTIC ACTION

External antiparasitic.

COMPOSITION

Each 100 mL of product contains:
Fipronil.....0.25 g
Excipients q.s.p.....100 mL

INDICATIONS

FIPROKILL[®] SPRAY is effective in the control and treatment of flea (*Ctenocephalides felis*) and tick (*Rhipicephalus sanguineus*) infestation in dogs and cats from 2 days of age, providing a rapid knock-down effect against adult fleas.

EFFECTIVENESS

FIPROKILL[®] SPRAY in dogs provides a protection period of up to 3 months against fleas and up to 5 weeks against ticks. In cats, the period of protection against fleas extends for 4 weeks.

MODE OF APPLICATION

1. Apply the product in ventilated environments and make sure to use gloves. Avoid contact with

- skin. If there is contact with hands, wash with plenty of water after handling the product.
2. Apply always against the grain, as indicated in the figure, from a distance of approximately 20 cm.
 3. The areas where the product should be applied are the belly and the flanks (chest, abdomen, armpits, shoulders and paws).
 4. For long coat animals, separate the hair and apply the product “against the grain”, until the root.
 5. Rub the coat for achieving the product penetrates the skin.
 6. After applying, do not expose the pet to heat sources or fire. Let dry naturally. Do not use hairdryer.
 7. For treating the face, pour a little of the product in the palm of the gloved hand, and then apply it rubbing gently the face, avoiding touching the eyes, mouth and mucous membranes.
 8. Long or thick coat animals may require the application of a larger quantity of product.
 9. It is recommended to not bathe the pet within 48 hours before and after applying Fiprokill® Spray.
 10. The product is safe in kittens and puppies from 2 days of age, provided that the application is carried out following the manufacturer’s instructions.

ROUTE OF ADMINISTRATION AND DOSAGE

Dose of active ingredient:

7.5 mg - 15 mg/kg of body weight.

Dosage of product:

- **Fiprokill Spray 50 mL and 100 mL bottle:** Apply FIPROKILL® Spray Solution, approximately 6 to 12 sprays by kg of body weight (wetting the hair root), in such a way the animal becomes totally wet.
- **Fiprokill Spray 250 mL bottle:** Apply FIPROKILL® Spray Solution, approximately 2 to 4 sprays by kg of body weight (wetting the hair root), in such a way the animal becomes totally wet.

Long or thick coat animals may require the application of a larger quantity of product.

CONTRAINDICATIONS

- Do not apply to rabbits.
- Do not apply in animals with known hypersensitivity to fipronil.

PRECAUTIONS

- Apply in ventilated environments.
- Wear gloves when applying the product.
- Do not smoke, eat or drink while handling and applying the product.
- Avoid contact with the treated animal until its fur is completely dry.
- Avoid sleeping with the recently treated animal.
- In case of contact with skin, eyes or mucous membranes, wash immediately with plenty of water.
- In case of accidental ingestion, go immediately to a medical center and show the product label.

WARNINGS

- Keep out of the reach of children.
- Administer with caution in weakened and senile animals.
- flammable product. Keep away from heat sources

ADVERSE EFFECTS

- If the animal licks itself, there may be a brief period of hypersalivation (mainly in cats).
- Transient skin reactions at the application site such as erythema, itching, or alopecia may occur.
- In the event of hypersensitivity or local irritation, it is recommended not to repeat the treatment.

OBSERVATIONS

Frequently asked questions:

What happen if my pet licks itself after the application? What would happen to my pet?

FIPROKILL® SPRAY is formulated with an active ingredient that is safe for mammals; however, after applying the product, avoid your pet licks itself until its coat is dry.

Can I let my children caress the animals after the application? How long should I wait for?

It is advisable the pet is totally dry before touching it again. Otherwise, wash the hands with plenty of water.

Can I bathe my pet without altering the efficacy of FIPROKILL® SPRAY?

It is recommended to not bathe your pet at least 48 hours after applying the product.

CONSERVATION

Store between 15 and 30 °C, protected from light. Once the container is opened, use within 12 weeks. Discard unused product after that time period

CONDITION OF SALE

OTC product (non-prescription)

PRESENTATION

Bottle with spray valve with 50 mL, 100 mL and 250 mL

PREPARED BY

Drag Pharma Laboratory

RECORDS

Chile: Reg. SAG N° 1872
Panamá: Reg. N° RF-4541-19
Rep. Dominicana: Reg. N° 8664
Bolivia: Reg. SENASAG PUV-F N° 007256/16
Perú: Registro SENASA F.87.30.I.0193
Uruguay: Reg. MGAP N° 2020A00783

COUNTRIES WHERE IT IS MARKETED

Imported and distributed in Bolivia by:

ZOOFARMA
TEL: +(591)222-3357
Street Díaz Romero 1339, La Paz.

Imported and Distributed in Peru by:

Representaciones Durand SAC.

Av. Manuel Olgúin No. 501 Office No. 604 Santiago de Surco Lima.

Importer in Uruguay: VIVAFIL S.A.

Av. 18 de Julio 1746 apt 601 - Montevideo - Uruguay.

grupotecnovet@gmail.com Technical Director: DMTV Diego Cuadrado.

NOTE

Special precautions for disposal of unused product or waste material: Dispose of this product with care together with household waste. Do not dispose of containers with product residues on the ground or in watercourses, as the product is toxic to birds, bees and fish. For expired or unused products, contact the manufacturing laboratory.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian