KET-10[®] BOVINE - Injectable solution

ANALGESIC, ANTIPYRETIC AND NON-STEROIDAL ANTI-INFLAMMATORY



In cattle, indicated as an antiinflammatory, analgesic and
antipyretic treatment, particularly in:
respiratory conditions, acute
bacterial mastitis, calf hyperthermia
and in musculoskeletal conditions
such as lameness, arthritis, trauma.
In pigs, indicated in the relief of pain,
fever and inflammation in respiratory
processes and in metritis syndrome,
mastitis, agalactia, in conjunction
with antibiotics.

Technical Specification

SPECIES

Bovine and pigs.

DOSAGE FORM

Injectable solution.

THERAPEUTIC ACTION

ANALGESIC, ANTIPYRETIC AND NON-STEROIDAL ANTI-INFLAMMATORY

COMPOSITION

Each mL of injectable solution contains:

Ketoprofen...... 100 mg Excipients q.s.p...... mL

INDICATIONS

In cattle, indicated as an anti-inflammatory, analgesic and antipyretic treatment, particularly in: respiratory conditions, acute bacterial mastitis, calf hyperthermia and in musculoskeletal conditions such as lameness, arthritis, trauma.

In pigs, indicated in the relief of pain, fever and inflammation in respiratory processes and in metritis syndrome, mastitis, agalactia, in conjunction with antibiotics.

ROUTE OD ADMINISTRATION AND DOSAGE

Bovine

Administration route: Intramuscular and intravenous

Dose: 3 mg of Ketoprofen / Kg of live weight, equivalent to 3 mL of product / 100 Kg of weight.

Pigs

Administration route: Intramuscular.

Dose: 3 mg of Ketoprofen / Kg of live weight, equivalent to 3 mL of product / 100 Kg of weight.

DRUG INTERACTIONS

Simultaneous use with other non-steroidal anti-inflammatory drugs can increase side effects.

CONTRAINDICATIONS

- Do not administer in pregnant females.
- Do not use in cases of severe kidney failure.
- Do not use simultaneously with diuretics or anticoagulants.
- Do not use in animals with heart or liver diseases, gastrointestinal ulcers or blood dyscrasias.
- Do not use in animals with known hypersensitivity to the active substance.

WARNINGS

Keep out of reach of children.

GUARD PERIOD

Meat: 4 days.
Milk: 0 days

CONSERVATION

Store at a temperature between 2 $^{\circ}$ and 30 $^{\circ}$ C, protected from light. Once opened, use the product within 3 months. Discard the excess product after that period of time.

CONDITION OF SALE

To be supply only on veterinary prescription.

PRESENTATION

50 mL vial

PREPARED BY

Drag Pharma Laboratory.

RECORDS

Chile: Reg. SAG N° 1472

Bolivia: Reg. SENASAG PUV-F N° 007258/16 Perú: Registro SENASA F.99.01.I.0069

COUNTRIES WHERE IT IS MARKETED

Imported and distributed in Bolivia by: AGROGUARANI SRL TEL: + (591) 314-1401 Santa Cruz de la Sierra, Bolivia Imported and Distributed in Peru by Representaciones Durand SAC. Av. Manuel Olguín No. 501 Office No. 604 Santiago de Surco Lima.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian