

PANVERMIC® Oral Tablet

BROAD SPECTRUM INTERNAL ANTIPARASITIC.



Broad spectrum internal antiparasitic in a single dose. Effective against nematodes and cestodes, such as: *Toxocara canis*, *Toxascaris leonina*, *Ancylostoma caninum*, *Hookworm stenocephala*, *Taenia hydatigena*, *Taenia pisiformis*, *Taenia taeniformis*, *Dipylidium caninum*, and *Echinococcus granulosus*.

Technical Specification

SPECIES

Dogs.

DOSAGE FORM

Oral Tablet

THERAPEUTIC ACTION

Broad spectrum internal antiparasitic.

COMPOSITION

Each tablet contains:

Praziquantel 50 mg
Mebendazole 220 mg
Pyrantel Pamoate 144 mg
(Equivalent to 50 mg of Pyrantel base)
Excipients c.s.p 1 tablet

INDICATIONS

Broad spectrum internal antiparasitic in a single dose. Effective against nematodes and cestodes, such as: *Toxocara canis*, *Toxascaris leonina*, *Ancylostoma caninum*, *Hookworm stenocephala*, *Taenia hydatigena*, *Taenia pisiformis*, *Taenia taeniformis*, *Dipylidium caninum*, and *Echinococcus granulosus*.

ROUTE OF ADMINISTRATION AND DOSAGE

Administer orally, directly into the oral cavity or mixed with food.

- Dose of active ingredients:

Administer 22 mg / Kg of Mebendazole; 5 mg / Kg of Praziquantel and 5 mg / Kg of Pyrantel as a base.

- Product dosage:

1 tablet for every 10 kg of weight. Single dose.

For the treatment of *Trichuris vulpis* infestations, administer for 3 consecutive days.

DRUG INTERACTIONS

- Do not administer together with Piperazine as it could antagonize the effect of Pyrantel Pamoate.
- Do not administer in conjunction with Morantel or Levamisole due to their mechanisms of action and toxicity similar to Pyrantel.

CONTRAINDICATIONS

- Do not administer in dogs with known hypersensitivity to the active ingredients.
- Do not administer to puppies under 4 weeks of age.
- Do not administer to dogs with liver failure, malnourished, dehydrated or anemic.

SPECIAL PRECAUTIONS FOR THE OPERATOR

- Wash your hands after administering the product.
- In case of accidental ingestion, do not induce vomiting. Get medical help.

WARNINGS

Keep out of the reach of children.

ADVERSE EFFECTS

In some animals they can present in a mild and transient way: nausea, vomiting, anorexia, abdominal pain and diarrhea.

OBSERVATIONS

Special precautions for disposal of unused product or waste material:

Discard any unused product remains in its original container. Do not throw the empty container or with product remains, in rivers, lakes or streams of natural water. Dispose of this product waste carefully with household waste.

CONSERVATION

Keep in a cool and dry place, at room temperature between 15° and 30° C.

CONDITION OF SALE

Sale with veterinary prescription.

PRESENTATION

Box with 2 tablets

PREPARED BY

Laboratorio Drag Pharma Chile Invetec S.A.

RECORDS

Reg. SAG N° 1689

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian