

PRAZIVERMIC® Oral tablet

BROAD SPECTRUM INTERNAL ANTIPARASITIC.



Broad spectrum anthelmintic, effective against cestodes and nematodes present in dogs, such as: *Toxocara canis*, *Ancylostoma caninum*, *Taenia hydatigena*, *Taenia pisiformis*, *Echinococcus granulosus*, and *Dipylidium caninum*.

Technical Specification

SPECIES

Dogs.

DOSAGE FORM

Oral tablet

THERAPEUTIC ACTION

Broad spectrum internal antiparasitic.

COMPOSITION

Each tablet contains:

Levamisole Hydrochloride 80 mg

Praziquantel 50 mg

Excipients q.s.p 1 tab.

INDICATIONS

Broad spectrum anthelmintic, effective against cestodes and nematodes present in dogs, such as: *Toxocara canis*, *Ancylostoma caninum*, *Taenia hydatigena*, *Taenia pisiformis*, *Echinococcus granulosus*, and *Dipylidium caninum*.

ROUTE OF ADMINISTRATION AND DOSAGE

Route of administration and method of use: Oral route.
Administer directly into the oral cavity or mixed with food.

Dose:

Administer 1 tablet for every 10 kilos of weight, equivalent to 8 mg / Kg of Levamisole Hydrochloride and 5 mg / Kg of Praziquantel, in a single dose.

DRUG INTERACTIONS

- It is recommended to avoid the administration of Levamisole in conjunction with Chloramphenicol, since deaths have been reported when they have been administered concomitantly.
- Cholinesterase inhibitor drugs (eg, organophosphates, Neostigmine) can potentiate the toxic effects of Levamisole.
- Nicotinic-type components (eg, Pyrantel, Morantel, Diethylcarbamazine) can enhance the toxic effects of Levamisole.
- Levamisole administered together with Warfarin increases the risk of bleeding.
- Interactions of Praziquantel with other medicinal products for veterinary use have not been described.

CONTRAINDICATIONS

- Do not administer to dogs under 4 weeks of age.
- Do not administer to patients with a history of gastric intolerance.
- Do not administer to animals with hypersensitivity to any of the active principles.

Use during pregnancy and lactation:

Do not administer to pregnant or lactating females.

SPECIAL PRECAUTIONS FOR THE OPERATOR

- Wash your hands with soap and plenty of water after handling the product.
- In the case of contact with the eyes, it is recommended to wash with plenty of water for 15 minutes. If irritation develops and persists, see doctor.
- In case of accidental ingestion, do not induce vomiting. Get medical help.

WARNINGS

Keep out of the reach of children.

ADVERSE EFFECTS

In some cases, hypersalivation, vomiting or muscle tremors may be observed, which are temporary symptoms and disappear spontaneously.

OBSERVATIONS

Special precautions for disposal of unused product or waste material:

Discard any unused product remains in its original container. Do not throw the empty container or with product remains, in rivers, lakes or streams of natural water. Do not reuse the container. Dispose of this product waste carefully with household waste.

CONSERVATION

Store in a cool, dry place, at room temperature between 15° and 30° C.

CONDITION OF SALE

Sale with veterinary prescription.

PRESENTATION

Box with 2 tablets

PREPARED BY

Laboratorio Drag Pharma Chile Invetec S.A.

RECORDS

Chile: Reg. SAG N° 0362

Perú: Registro SENASA F.08.21.I.0925

COUNTRIES WHERE IT IS MARKETED

Imported and Distributed in Peru by:
Representaciones Durand SAC.
Av. Manuel Olguín No. 501 Office No. 604 Santiago de Surco Lima.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian