

ROSTRUM[®] PLUS - Otic Emulsion

ANTIBACTERIAL AND ANTIFUNGAL



Indicated for the treatment of acute or chronic external otitis in dogs, caused by bacteria and fungi sensitive to Enrofloxacin and / or Silver Sulfadiazine.

Technical Specification

SPECIES

Dogs.

DOSAGE FORM

Otic, antibacterial and antifungal emulsion

THERAPEUTIC ACTION

Antibacterial and antifungal

COMPOSITION

Each mL of emulsion contains:
Enrofloxacin 5 mg
Silver Sulfadiazine 10 mg
Excipients q.s.p 1 mL

INDICATIONS

Indicated for the treatment of acute or chronic external otitis in dogs, caused by bacteria and fungi sensitive to Enrofloxacin and / or Silver Sulfadiazine.

MODE OF APPLICATION

Clean the ear canal, then administer Rostrum[®] Plus in sufficient quantity to cover the external ear canal, gently massaging the ear to ensure a complete and even distribution of the medicine throughout the canal.

ROUTE OF ADMINISTRATION AND DOSAGE

Topical administration.

Product dosage: 5 to 10 drops in dogs weighing 15 kilos or less, and 10 to 15 drops in dogs weighing more than 15 kilos. Administer 2 times a day for 14 days.

DRUG INTERACTIONS

It is not recommended to use concomitantly with other pharmaceutical products.

CONTRAINDICATIONS

- Do not use in dogs with hypersensitivity to Quinolones and / or Sulfonamides.
- Do not use in dogs with perforated tympanic membrane.
- Do not use during pregnancy and lactation.

SPECIAL PRECAUTIONS FOR USE

- The use of this combination has not been evaluated in dogs with perforation of the tympanic membrane. Consequently, before administering the product, the external auditory canal should be examined in order to ensure that there is no rupture of the tympanic membrane, in order to avoid contamination of the middle ear, as well as vestibular and cochlear lesions. If during the administration of the product there are signs of deafness or vestibular alterations, discontinue its use.
- Quinolones should be used with caution in animals with central nervous system (CNS) disorders. In these animals, in rare cases quinolones have been associated with CNS stimulation, which can generate seizures.
- Fluoroquinolones, such as Enrofloxacin, should not be used as the first line of treatment, unless there is no therapeutic alternative available. When used as a second treatment, it should be based on susceptibility studies.

SPECIAL PRECAUTIONS FOR THE OPERATOR

Wash your hands after administering the product.

WARNINGS

Keep out of the reach of children.

ADVERSE EFFECTS

- With the exception of possible cartilage abnormalities in young animals, the adverse effect profile of Enrofloxacin is generally limited to gastrointestinal problems (vomiting, anorexia), although rare incidents of elevated liver enzymes, ataxia, seizures, depression have also been reported. and lethargy.
- No evidence of systemic toxicity due to Silver Sulfadiazine has been reported in dogs, even 1% Silver Sulfadiazine ear suspensions have been used in dogs for more than three months without incident.

OBSERVATIONS

Shake before using.

Special precautions for disposal of unused product or waste material:

Dispose of any unused product remains in its original, tightly closed container. Dispose of this product waste carefully with household waste.

CONSERVATION

Keep in a cool place, at a temperature between 15° and 30° C, protected from light.

CONDITION OF SALE

Sale with veterinary prescription.

PRESENTATION

Dropper bottle with 15 mL

PREPARED BY

Laboratorio Drag Pharma Chile Invetec S.A.

RECORDS

Chile: Reg. SAG N° 1523
Bolivia: Reg. SENASAG PUV-F-N° 006147/14
El Salvador: VE2014084963

COUNTRIES WHERE IT IS MARKETING

Imported and distributed in Bolivia by:
ZOOFARMA

TEL: +(591)222-3357
Street Díaz Romero 1339, La Paz.

Distributed in El Salvador by:
Rafael Alfredo Alfaro Castillo.

Imported and Distributed in Peru by:
Representaciones Durand SAC.
Av. Manuel Olguín No. 501 Office No. 604 Santiago de Surco Lima.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian