

TOPFENICOL® Injectable solution

ANTIMICROBIAL FOR CATTLE AND PIGS.



In cattle, it is indicated in bovine respiratory disease, in which susceptible strains of *Pasteurella haemolytica*, *Pasteurella multocida*, *Haemophilus somnus*, *Actinomyces pyogenes* are associated. Its therapeutic action is also recommended in infectious keratoconjunctivitis caused by *Moraxella bovis*.

In pigs, its use is related to respiratory disease. The strains causing the syndrome that are described as susceptible are *Pasteurella multocida*, *Mycoplasma hyopneumoniae*, *Salmonella choleraesuis*, *Mycoplasma hyorhinus*, *Bordetella bronchiseptica*, *Streptococcus suis*.

Technical Specification

SPECIES

Bovines and pigs.

DOSAGE FORM

Injectable solution

THERAPEUTIC ACTION

Antimicrobial for cattle and pigs.

COMPOSITION

Each 1 mL of product contains:
Florfenicol 300 mg
Excipients q.s.p 1 mL

PROPERTIES

Florfenicol is a long-acting injectable antibiotic with action on a wide variety of aerobic and anaerobic Gram (+) and Gram (-) bacteria, isolated from cattle and pigs.

INDICATIONS

In cattle, it is indicated in bovine respiratory disease, in which susceptible strains of *Pasteurella haemolytica*, *Pasteurella multocida*, *Haemophilus somnus*, *Actinomyces pyogenes* are associated. Its therapeutic action is also recommended in infectious keratoconjunctivitis caused by *Moraxella bovis*.

In pigs, its use is related to respiratory disease. The strains causing the syndrome that are described as susceptible are *Pasteurella multocida*, *Mycoplasma hyopneumoniae*, *Salmonella choleraesuis*, *Mycoplasma hyorhinus*, *Bordetella bronchiseptica*, *Streptococcus suis*.

ROUTE OF ADMINISTRATION AND DOSAGE

Dose of the active ingredient:

- Cattle: 20 mg / Kg of live weight, intramuscularly.
- Pigs: 15 mg / Kg of live weight, intramuscularly.

Repeat a second dose 48 hours after starting treatment.

Product dosage:

- Cattle: 6.7 mL / 100 Kg of weight, intramuscularly.
- Pigs: 0.5 mL / 10 Kg of live weight, intramuscularly.

Repeat a second dose 48 hours after starting treatment.

CONTRAINDICATIONS

- Do not administer to animals that have previously shown hypersensitivity to chloramphenicol or florfenicol.
- Do not use in cattle intended for encasing.
- Do not use on bristles intended for setting.
- Do not use in pregnant or lactating females.

SPECIAL PRECAUTIONS FOR USE

If signs of respiratory disease persist 48 hours after the second dose, reevaluate the diagnosis.

ADVERSE EFFECTS

In cattle, a temporary drop in food and water consumption may occur at the start of therapy.
In pigs, transient perianal inflammation may occur at the start of treatment.

OBSERVATIONS

Employment during pregnancy and lactation:

Do not use in animals intended for reproduction, or in pregnant or lactating females. Efficacy period: 36 months

Efficacy period

36 months

Disposal of waste material:

Discard any unused product remains in its original container. Do not throw the empty container or with the rest of the product in rivers, lakes or streams of water. Dispose of the remains of this product with caution together with household waste.

CONSERVATION

Store between 15 and 30 ° C, in a cool and dry environment, protected from light and in its original container.

CONDITION OF SALE

To be supply only on veterinary prescription.

PRESENTATION

100 mL vial

PREPARED BY

Laboratorio Drag Pharma Chile Invetec.S.A.

RECORDS

Chile: Reg. SAG N° 2022

Bolivia: Reg. SENASAG PUV-F nº 007492/16

Perú: Registro SENASA F.03.01.I.1844

COUNTRIES WHERE IT IS MARKETED

Imported and distributed in Bolivia by:

ZOOFARMA

TEL: +(591)222-3357

Street Díaz Romero 1339, La Paz.

Imported and Distributed in Peru by Representaciones Durand SAC.

Av. Manuel Olgún No. 501 Office No. 604 Santiago de Surco Lima.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian