CABATINA[®] Oral solution

ANTI-HAEMORRHAGIC ANTI-HAEMORRHAGIC FOR ANTICOAGULANT RODENTICIDE POISONING.



Cabatina[®] is a product indicated for the treatment of intoxications generated by anticoagulant rodenticides and hemorrhagic processes related to the defective formation of vitamin K-dependent coagulation factors.

Technical Specification

SPECIES

Dogs and cats.

DOSAGE FORM

Oral solution

THERAPEUTIC ACTION

Anti-haemorrhagic Anti-haemorrhagic for anticoagulant rodenticide poisoning.

COMPOSITION

Each 1 mL of product contains: Vitamin K1 (Phytomenadione) 20 mg Excipients c.s.p 1mL

PROPERTIES

Vitamin K is a cofactor in the last stage of the hepatic synthesis of coagulation factors II (Prothrombin), VII (Proconvertin), IX (thromboplastin component of plasma) and X (Stuart factor). The process of formation of vitamin K-dependent clotting factors can be interrupted for various reasons, such as congenital diseases, vitamin K deficiency as a result of low dietary intake, malabsorption syndromes, broad-spectrum antibiotic therapies, or others.

In veterinary medicine, the most frequent cause of the cessation of K-dependent coagulation factor formation is intoxication by anticoagulant rodenticides. Vitamin K1 antagonizes the effects of anticoagulant rodenticidal agents and allows to reactivate the production process of K-dependent coagulation factors.

INDICATIONS

Cabatina[®] is a product indicated for the treatment of intoxications generated by anticoagulant rodenticides and hemorrhagic processes related to the defective formation of vitamin K-dependent coagulation factors.

ROUTE OD ADMINISTRATION AND DOSAGE

Oral.

Dosage:

Dose of the active principle:

Dogs: 2 mg / Kg of vitamin K1, once a day, for 3 weeks. Cats: 5 mg / Kg of vitamin K1, once a day, for 3 weeks.

Product dosage:

Dogs: 1 mL of Cabatina[®] for every 10 Kg of body weight, once a day, for 3 weeks. Cats: 1 mL Cabatina[®] for every 4 Kg of body weight, once a day, for 3 weeks.

Recommendations:

To improve its absorption, it is recommended to administer Cabatina® together with a tablespoon of a food rich in fatty acids, such as canned food for dogs or cats.

DRUG INTERACTIONS

The following medications can prolong or potentiate the effects of anticoagulants and antagonize some of the therapeutic actions of Vitamin K1: Phenylbutazone, Aspirin, Chloramphenicol, Sulfonamides (including Sulfa / Trimethoprim), Diazoxide, Allopurinol, Cimetidine, Metronidazole, anabolic steroids, Erythromycin, Ketoconazole, Propanolol and thyroid drugs. The joint administration of Vaseline can decrease the absorption of Vitamin K.

CONTRAINDICATIONS

- Do not administer to animals with known hypersensitivity to vitamin K.
- Do not administer in animals with severe liver failure.

PRECAUTIONS

This product does not correct hypoprothrombinemia resulting from hepatocellular damage.

SPECIAL PRECAUTIONS FOR THE OPERATOR

- In case of contact with the skin, it is recommended to wash hands with soap and plenty of water. If irritation develops and persists, see doctor.
- In the case of contact with the eyes, it is recommended to wash with plenty of water for 15 minutes. If irritation develops and persists, see doctor.
- In case of ingestion, do not induce vomiting. Get medical help.

WARNINGS

Keep out of the reach of children.

OBSERVATIONS

Special precautions for disposal of unused product or waste material:

Discard any unused product remains in its original container. Dispose of the waste of this product with care together with household waste. Contact the manufacturing company or companies

specialized in waste disposal, to receive recommendations on the disposal of expired or unused products.

CONSERVATION

- Store at a temperature between 15 and 30 ° C, protected from light.
- Once opened, use within 8 weeks. Discard the unused product after that period of time.

CONDITION OF SALE

Sale with Veterinary Medical prescription.

PRESENTATION

25 mL bottle

PREPARED BY

Laboratorio Drag Pharma Chile Invetec S.A.

RECORDS

Chile: SAG Reg. No. 2096 Peru: Reg. SENASA F.77.32.1.0053 Imported and distributed by Representaciones Durand SAC. Av. Manuel Olguín N° 501 Office N° 604 Santiago de Surco Lima. Bolivia: Reg. SENASAG PUV-No 009805/21 Imported and distributed by: ZOO PHARMA VETERINARY INPUTS S.R.L. Diaz Romero N° 1339 Miraflores Zone, La Paz-Bolivia Telephone: 591 2223357 Costa Rica: Reg. No. MV-7129

COUNTRIES WHERE IT IS MARKETED

Imported and distributed in Bolivia by:

ZOOFARMA TEL: +(591)222-3357 Street Díaz Romero 1339, La Paz.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian