

# EQUIMIC® Oral Gel

## INTERNAL ANTIPARASITIC.



Single dose broad spectrum internal antiparasitic; effective against adult, migratory and / or larval stages of gastrointestinal parasites such as: *Strongylus vulgaris*, *Strongylus equinus*, *Strongylus edentatus*, *Oxyuris equi*, *Parascaris equorum*, small strongyles, *Strongyloides westeri*, *Trichostrongylus axei*, *Habronema spp.* and *Gasterophilus spp.*

## Technical Specification

### SPECIES

Horses.

### DESCRIPTION

Broad spectrum internal antiparasitic for horses

### DOSAGE FORM

Oral gel.

### THERAPEUTIC ACTION

Internal Antiparasitic.

### COMPOSITION

Each 1 mL of gel contains:  
Ivermectin.....14.0 mg  
Excipients q.s.p.....1 mL

### INDICATIONS

Single dose broad spectrum internal antiparasitic; effective against adult, migratory and / or larval stages of gastrointestinal parasites such as: *Strongylus vulgaris*, *Strongylus equinus*, *Strongylus edentatus*, *Oxyuris equi*, *Parascaris equorum*, small strongyles, *Strongyloides westeri*, *Trichostrongylus axei*, *Habronema spp.* and *Gasterophilus spp.*

### MODE OF APPLICATION

- **Foals:** Start the treatment between 6-8 weeks of age, repeating the treatment at intervals of 6 to 8 weeks.

- **Mares:** It is recommended to treat 1 month before the due date and repeat 60 days after delivery.
- **Adult horses:** Treatment every 90 days. The repetition of the treatment depends on the parasite load and the season of the year.

#### ROUTE OF ADMINISTRATION AND DOSAGE

---

**Route of administration:** Oral.

**Active ingredient dose:**  
0.2 mg/Kg of weight, single dose.

**Product dose:**  
1 ml/70 kg of weight (0.2 mg/Kg), single dose.

Each syringe reaches 700 kg of weight.

#### CONTRAINDICATIONS

---

- Do not use in foals under 4 weeks of age.
- Do not administer to horses with known hypersensitivity to Ivermectin.

#### SPECIAL PRECAUTIONS FOR THE OPERATOR

---

- In case of accidental ingestion, do not induce vomiting. Consult a doctor as soon as possible.
- In case of contact with skin or eyes, wash with plenty of water. If irritation occurs, consult a doctor.
- Wash hands after administration of the product.

#### WARNINGS

---

- Keep out of reach of children.

#### ADVERSE EFFECTS

---

Adverse effects are observed in a low percentage of animals to which oral and injectable formulations containing Ivermectin are administered simultaneously.

In cases of severe infestations, signs of transient colic may be observed.

#### GUARD PERIOD

---

Do not administer to horses whose meat is intended for human consumption.

#### OBSERVATIONS

---

Pleasant apple flavor.

#### **Special precautions for disposal of unused product or waste material:**

Discard any unused product remains in its original container. Do not throw the empty container or with product remains, in rivers, lakes or streams of water. Dispose of the waste of this product with care together with household waste.

#### CONSERVATION

---

Store in a cool, dry place, between 15 and 30 ° C.  
Use immediately once opened and discard the excess product.

#### CONDITION OF SALE

---

To be supply only on veterinary prescription.

#### PRESENTATION

---

10 mL graduated syringe.

#### PREPARED BY

---

Drag Pharma Laboratory.

#### RECORDS

---

Chile: Reg. SAG No. 1655-B

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian