

FEBRECTAL[®] EQUINE - Injectable solution

ANTI-INFLAMMATORY, ANALGESIC AND ANTIPYRETIC.



Anti-inflammatory, analgesic and antipyretic. Recommended to relieve inflammation, fever and pain associated with musculoskeletal disorders and relieve pain associated with colic.

Technical Specification

SPECIES

Horses, cattle and pigs.

DOSAGE FORM

Injectable solution.

THERAPEUTIC ACTION

Anti-inflammatory, analgesic and antipyretic.

COMPOSITION

Each mL of solution contains:

Flunixin Meglumine.....83 mg

(Equivalent to 50 mg of Flunixin)

Excipients q.s.p.....1 mL

INDICATIONS

Anti-inflammatory, analgesic and antipyretic. Recommended to relieve inflammation, fever and pain associated with musculoskeletal disorders and relieve pain associated with colic.

ROUTE OF ADMINISTRATION AND DOSAGE

- Horses: 1.1 mg/Kg/day (1 mL/45 Kg of weight) EV or IM, repeat for a maximum of 5 days.
- Cattle: 2.2 mg/Kg/day (2 mL/45 Kg of weight) IV, in one or two administrations.
- Pigs: 2.2 mg/Kg/day (2 mL/45 Kg of weight) IM for a maximum of 3 days.

DRUG INTERACTIONS

- Concurrent administration of potentially nephrotoxic drugs should be avoided. - Flunixin can reduce the renal excretion of some drugs, increasing their toxicity. - The drug should not be administered together with other NSAIDs or glucocorticoids, as it can increase the toxicity of both, especially at the gastrointestinal level, increasing the risk of suffering from gastrointestinal ulcers.

CONTRAINDICATIONS

- Do not use in animals with hypersensitivity to the active ingredient or other NSAIDs.
- Do not use in animals suffering from heart, liver or kidney disease.
- Do not use in animals with lesions of the gastrointestinal tract, such as ulcers and hemorrhages.
- Do not use when there are signs of blood dyscrasias or impaired hemostasis.
- Do not use in cases of colic caused by ileus and associated with dehydration.
- Do not use in animals suffering from chronic musculoskeletal disorders.
- Do not administer to animals under 72 hours of life.
- Do not administer to pregnant or lactating females or to breeding animals.

PRECAUTIONS

Special precautions for disposal of unused product or waste material:
Empty containers can be discarded as domestic waste, without any special precautions. Nope
Dispose of containers with product residues on the ground or in watercourses. For products expired
or not
used contact the manufacturing laboratory

SPECIAL PRECAUTIONS FOR THE OPERATOR

- Do not eat, drink or smoke while handling the product.
- Avoid self-injection.

WARNINGS

Keep out of reach of children.

ADVERSE EFFECTS

- In horses after IM injection, cases of localized inflammation have been reported, induration, stiffness and sweating.
- Accidental intra-arterial injection may cause stimulation of the central nervous system with signs such as ataxia, hyperventilation and muscle weakness. Clinical signs are transient and They generally do not require any treatment.
- Flunixin is a relatively safe agent for use in the horse, but there is the possibility of Gastrointestinal intolerance, hypoproteinemia, and hematologic abnormalities may occur.
- In horses and cattle, unusual anaphylactic reactions have been reported, mainly after rapid IV administration.
- Hematochezia and hematuria have been reported in cattle treated longer than 3 day recommendation.
- In pigs, no adverse reactions have been reported, except for some cases of irritation at the point of injection.

GUARD PERIOD

- Do not use in horses intended to human consumption.
- Bovine: 7 days for meat, 2 days for milk.
- Pigs: 21 days for meat.

CONSERVATION

Store between 15° and 30°C, protected from light. Once the container is opened, use within 28 days. Discard unused product after that time period.

CONDITION OF SALE

To be supply only on veterinary prescription.

PRESENTATION

20 and 50 mL ampule.

PREPARED BY

Drag Pharma Laboratory.

RECORDS

- Chile: Reg. SAG No. 393
- Dominican Republic: Reg. No. 5602

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian