

FIPRODRAG® 4,02 mL - Dogs between 40 kg and 60 kg of body weight

EXTERNAL ANTIPARASITIC.



FIPRODRAG® is indicated for the treatment and control of fleas (*Ctenocephalides felis* and *Ctenocephalides canis*), lice (*Trichodectes canis*) and ticks (*Rhipicephalus sanguineus*) infestation in dogs over 10 weeks old. It is also recommended as adjuvant therapy in flea allergy dermatitis (FAD) in dogs. In this case, it is recommended a monthly application of product, including the animals cohabiting with them.

Efficacy: With a single application, FIPRODRAG® allows to control reinfestation in dogs, during a period of at least 8 weeks against fleas, and 5 weeks against ticks. If parasite burden is high, a monthly application of product is advisable. For its lipophilic properties, FIPRODRAG® resists baths carried out 48 hours after its application. Do not bathe your pet 48 hours before and after applying the product.

Technical Specification

SPECIES

Dogs between 40 kg and 60 kg of body weight.

DOSAGE FORM

Topical solution.

THERAPEUTIC ACTION

External antiparasitic.

COMPOSITION

Each mL of solution contains:

Fipronil.....100 mg

Excipients q.s.p.....1 mL

PROPERTIES

Effective action against fleas, lice and ticks in dogs.

INDICATIONS

FIPRODRAG® is indicated for the treatment and control of fleas (*Ctenocephalides felis* and *Ctenocephalides canis*), lice (*Trichodectes canis*) and ticks (*Rhipicephalus sanguineus*) infestation in dogs over 10 weeks old. It is also recommended as adjuvant therapy in flea allergy dermatitis (FAD) in dogs. In this case, it is recommended a monthly application of product, including the animals cohabiting with them.

Efficacy: With a single application, FIPRODRAG® allows to control reinfestation in dogs, during a period of at least 8 weeks against fleas, and 5 weeks against ticks. If parasite burden is high, a monthly application of product is advisable. For its lipophilic properties, FIPRODRAG® resists baths carried out 48 hours after its application. Do not bathe your pet 48 hours before and after applying the product.

MODE OF APPLICATION

1. Remove the upper marked tip of the pipette.
2. Put the end of the pipette directly on the animal skin, at scapulae (shoulders) level or at the base of neck, trying to separate the hair.
3. Press and pour the full content of pipette on 2 or 3 points, as indicated in the figure.

ROUTE OF ADMINISTRATION AND DOSAGE

For external use, through skin application.

The dose of Fipronil formulated as a topical solution for dogs and cats has been established between 6.7 and 13.4 mg/kg body weight of the animal to be treated. In dogs between 40 kg and 60 kg of body weight, apply a pipette of 4.02 mL

CONTRAINDICATIONS

- Do not use on puppies less than 10 weeks old.
- Do not administer in rabbits.

PRECAUTIONS

- Administer with caution in weakened animals.
- The Bichon Frisé breed is particularly sensitive to side effects of Fipronil.

SPECIAL PRECAUTIONS FOR THE OPERATOR

Precautions to be taken by the person administering the product:

- Avoid touching the pipette content directly with your fingers.
- Wash your hands with soap and water after applying and handling the product.
- Do not eat food, smoke, or drink beverages during the application of product.
- Avoid handling your pet immediately after the treatment, until the coat is completely dry.
- Discard in a safe way the empty pipettes. Do not dispose the empty pipettes into natural waterways.

WARNINGS

- Keep out of reach of children.

- For external use only.
- The product may be harmful if eaten.
- The product is toxic to fishes, birds and bees.
- Flammable product.

SIDE EFFECTS

If the animal licks itself, a short period of hypersalivation may occur. If there is hypersensitivity or local irritation, it is recommended to not repeat the treatment.

OBSERVATIONS

- POISON.
- Flammable.

CONSERVATION

- Store in the original pack, at room temperature between 15 and 30°C, and in a fresh place.
- Keep out of heat sources.
- Do not store with food products.

CONDITION OF SALE

OTC product (non-prescription).

PRESENTATION

Box or Thermo-contracting Display containing 1 pipette of 4.02 mL for dogs from 40 to 60 kg of body weight.

PREPARED BY

Drag Pharma Laboratory.

RECORDS

SAG Reg. No. 2162

COUNTRIES WHERE IT IS MARKETED

Imported and distributed in Bolivia by:

ZOOFARMA
TEL: +(591)222-3357
Street Díaz Romero 1339, La Paz.

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian