

# INVECTINA® PLUS - Injectable Solution

## ENDECTOCIDE AND FASCIOLICIDE.



Indicated for the treatment and control of internal and external parasitism in cattle, sheep and goats.

Effective against mature and immature forms of gastrointestinal and lung nematodes, *Fasciola hepatica*, scabies, lice and *Oestrus ovis*.

## Technical Specification

### SPECIES

Bovine, sheep and goats.

### DOSAGE FORM

Injectable solution.

### THERAPEUTIC ACTION

Endectocide and fasciolicide.

### COMPOSITION

Each 1 mL of injectable solution contains:

Ivermectin.....5 mg

Closantel.....125 mg

Excipients q.s.p.....1 mL

### INDICATIONS

Indicated for the treatment and control of internal and external parasitism in cattle, sheep and goats. Effective against mature and immature forms of gastrointestinal and lung nematodes, *Fasciola hepatica*, scabies, lice and *Oestrus ovis*.

### ACTION SPECTRUM

**Bovine:** *Ostertagia ostertagi*, *Dictyocaulus viviparus*, *Trichostrongylus axei*, *T. columbriformis*, *Nematodirus helvetianus*, *Haemonchus contortus*, *Trichuris* spp. *Oesophagostomum radiatum*, *Cooperia oncophora*, *C. punctata*, *Fasciola hepatica*, *Linognathus* sp., *Psoroptes ovis*, *Sarcoptes scabiei*, *Haematopinus* sp.

**Sheep:** *Ostertagia circumcincta*, *O. trifurcata*, *Trichostrongylus columbriformis*, *T. vitrinus*, *T.*

*capricola*, *Nematodirus flicollis*, *N. spathiger*, *Haemonchus contortus*, *Cooperia curticei*, *Dictyocaulus sp.*, *Fasciola hepatica*, *Linognathus sp.*, *Oestrus ovis*, *Psoroptes sp.*, *Sarcoptes scabiei*.

**Goats:** *Ostertagia sp.*, *Trichostrongylus sp.*, *Nematodirus sp.*, *Haemonchus sp.*, *Cooperia sp.*, *Dictyocaulus sp.*, *Fasciola hepatica*, *Linognathus sp.*, *Oestrus ovis*, *Psoroptes sp.*, *Sarcoptes scabiei*.

#### ROUTE OF ADMINISTRATION AND DOSAGE

---

**Administration route:** Subcutaneous, single dose.

**Active ingredients dose:**

0.2 mg/Kg Ivermectin and 5 mg/Kg de Closantel.

**Product dose:**

Bovine, sheep and goats: 1 ml/125 kg of body weight, subcutaneous route, single dose.

#### CONTRAINDICATIONS

---

- Do not administer in animals less than 4 weeks of age.
- Do not administer to animals hypersensitive to avermectins, or to Closantel.

#### SPECIAL PRECAUTIONS FOR USE

---

- In case of individual hypersensitivity, suspend treatment.
- Do not administer intramuscularly or intravenously.
- For heavier animals, the dose should be divided into different injection sites to avoid discomfort in the area.

#### SPECIAL PRECAUTIONS FOR THE OPERATOR

---

- Do not handle by people who are hypersensitive to Ivermectin or Closantel.
- Use of gloves when handling the product.
- In case of contact with skin, eyes or mucous membranes, wash immediately with plenty of water. If there is irritation, go immediately to a medical center and show the product label.
- In case of accidental injection, go immediately to a medical center and show the product label.

#### WARNINGS

---

Keep out of reach of children.

#### GUARD PERIOD

---

**Meat:** 49 days.

**Milk:** Do not use the milk of treated animals for human consumption.

#### OBSERVATIONS

---

**Special precautions for disposal of unused product or waste material:**

Empty containers can be disposed of as household waste, without any special precautions. Do not dispose of containers with product remains on the ground or water courses. For expired or unused products, contact the manufacturing laboratory.

#### CONSERVATION

---

Store between 15 and 30 ° C, protected from light.

#### CONDITION OF SALE

---

To be supply only on veterinary prescription.

**PRESENTATION**

---

100 mL vial and 500 mL vial

**PREPARED BY**

---

Drag Pharma Laboratory.

**RECORDS**

---

- Chile: Reg. SAG No. 1375-B
- Dominican Republic: Reg. No. 5611

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian