

# INVETROID® Emulsifiable solution

---

EXTERNAL ANTIPARASITIC WITH RESIDUAL EFFECT.



Invetroid® is an effective antiparasitic in the control of lice (*Bovicola ovis*, *Bovicola bovis*, *Linognathus pedalis*, *Linognathus ovillus*, *Linognathus vituli*, *Haematopinus eurysternus*, *Haematopinus suis*), scabies mites (*Psoroptes bovis*, *Psoroptes bobiei*, *Psoroptes bobiei* var. , *Sarcoptes scabiei* var. *Suis*) and melophagi (*Melophagus ovinus*) in cattle, sheep and pigs.

## Technical Specification

### SPECIES

---

Bovine, sheep and pigs.

### DOSAGE FORM

---

Emulsifiable solution.

### THERAPEUTIC ACTION

---

External antiparasitic with residual effect.

### COMPOSITION

---

Each 100 mL of concentrated solution contains:

Cypermethrin..... 20 g

Excipients q.s.p.....100 mL

### INDICATIONS

---

Invetroid® is an effective antiparasitic in the control of lice (*Bovicola ovis*, *Bovicola bovis*, *Linognathus pedalis*, *Linognathus ovillus*, *Linognathus vituli*, *Haematopinus eurysternus*, *Haematopinus suis*), scabies mites (*Psoroptes bovis*, *Psoroptes bobiei*, *Psoroptes bobiei* var. , *Sarcoptes scabiei* var. *Suis*) and melophagi (*Melophagus ovinus*) in cattle, sheep and pigs.

### ROUTE OF ADMINISTRATION AND DOSAGE

---

#### Cattle and Sheep:

- **Baths for scabies and lice:** Dilute 1 liter of Invetroid® in 1,000 liters of water (equivalent to a final bath concentration of 0.02%). For replacement and reinforcement, use a dilution of 1.5 liters of the product in 1,000 liters of water (equivalent to a concentration of 0.03%); This is when the bathtub level has dropped by around 10%.

**- For dry replacement and reinforcement:** Dilute 500 mL of Invetroid® in 1,000 liters of water (equivalent to a final concentration of 0.01%). This dilution can be used as a spray, applying at a rate of 3-4 liters per animal.

**Sheep Melophagus (Melophagus ovinus: "false tick"):**

**- Immersion bath:** Loading: 1 liter of Invetroid® in 10,000 liters of water (equivalent to a final bath concentration of 0.002%). Replacement: 1 liter in 8,000 liters of water (equivalent to a concentration of 0.025%), when the level has dropped by 20%. It must be ensured that the animals are well wet, totally submerging them at least twice. Every 10,000 animals the bathroom must be renewed. Once the job is finished, all the remaining liquid must be removed.

**- Aspersión:** 1 liter of Invetroid® in 5,000 liters of water (equivalent to a final concentration of 0.004%). This system is recommended as a way to lower the melophaga load in freshly shorn animals.

**Pigs:**

**- Scabies and lice: Aspersión:** Dilute 1 mL of Invetroid® in 5 liters of water (equivalent to a final concentration of 0.004%). Apply at a rate of 3-4 liters per animal. Apply using a spray pump, distributing the product evenly over the animal.

**Frequency and duration of treatment:** Single dose.

**DRUG INTERACTIONS**

---

Do not use concomitantly with other pharmaceutical products.

**CONTRAINDICATIONS**

---

- Do not bathe in hours of intense heat, or when the animals are very tired and thirsty.
- Do not use during advanced stages of pregnancy or during lactation.
- Do not administer to animals with hypersensitivity to cypermethrin.

**SPECIAL PRECAUTIONS FOR USE**

---

- It must be ensured that both the immersion baths and the spray pumps are completely free of residues of other products that may have been applied previously.
- Use the mixture in a well-ventilated environment, avoiding contact with the animal's eyes.
- In case of rain threat, suspend the bath. In case of rain within 24 hours after bathing, repeat the treatment.
- Keep out of the reach of children.

**SPECIAL PRECAUTIONS FOR THE OPERATOR**

---

- The applicator must wear a mask and gloves as a protection element for handling the product.
- Avoid contact with the eyes, skin and mucous membranes, Cypermethrin is an irritant and can cause skin sensitization.
- Contact with treated animals should be avoided until they are completely dry.
- Wash contaminated clothing before reuse.
- In case of contact with the skin, wash with plenty of water. In case of eye contact, wash immediately with plenty of water for at least 15 minutes.

**ADVERSE EFFECTS**

---

In animals with hypersensitivity to cypermethrin, dermal irritation may be observed.

**GUARD PERIOD**

---

**Meat:**

- Bovine: 14 days
- Sheep and pigs: 7 days.

**Milk:** 7 days.

#### OBSERVATIONS

---

**Special precautions for disposal of unused product or waste material:**

Do not dispose of the empty container and / or product remains on the ground, rivers, lakes or streams of water. Any unused medicine or waste derived from it must be disposed of in an environmentally safe way.

#### CONSERVATION

---

Store below 30°C. Do not refrigerate or freeze.

Use immediately after opening and discard the excess product.

Use immediately once the solution is prepared and discard the excess product.

#### CONDITION OF SALE

---

To be supply only on veterinary prescription.

#### PRESENTATION

---

Bottle with 100 ml and 1 L.

#### PREPARED BY

---

Drag Pharma Laboratory.

#### RECORDS

---

**Chile:** Reg. SAG N° 955

**Rep. Dominicana:** N° 9685

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian