

# KAUPOL<sup>®</sup> BOVINE - Injectable solution

## VITAMINS, GLUCOSE AND MINERALS.



Indicated in dehydration and in glucose and mineral deficiencies, which can be caused by: bleeding, diarrhea, burns, postpartum deficiencies and / or poisoning.

In production animals Kaupol<sup>®</sup>, administered concomitantly with vitamins B<sub>1</sub>, B<sub>6</sub> and B<sub>12</sub>, is indicated as a hepatic, lipotropic and hematopoietic protector.

Especially recommended for animals that are weakened, convalescing, sick or subjected to long demands.

## Technical Specification

### SPECIES

Bovine, pigs, sheep and goats.

### DOSAGE FORM

Injectable solution.

### THERAPEUTIC ACTION

Vitamins, Glucose and Minerals.

### COMPOSITION

100 mL of injectable solution contains:

Riboflavin 5 Sodium phosphate.....	10 mg
Nicotinamide.....	200 mg
Anhydrous D-glucose (Dextrose).....	10.0 g
Sodium.....	11.96 mEq
Potassium.....	0.54 mEq
Magnesium.....	0.049 mEq
Calcium.....	0.163 mEq
Chloride.....	12.92 mEq
Excipients q.s.p.....	100 mL

### INDICATIONS

Indicated in dehydration and in glucose and mineral deficiencies, which can be caused by: bleeding, diarrhea, burns, postpartum deficiencies and / or poisoning.

In production animals Kaupol<sup>®</sup>, administered concomitantly with vitamins B<sub>1</sub>, B<sub>6</sub> and B<sub>12</sub>, is indicated as a hepatic, lipotropic and hematopoietic protector.

Especially recommended for animals that are weakened, convalescing, sick or subjected to long demands.

## ROUTE OF ADMINISTRATION AND DOSAGE

---

Slow and subcutaneous intravenous administration.

### Product dosage:

- Cattle and horses (greater than or equal to 400 kilos): 1000 - 2000 mL via E.V.
- Cattle (greater than or equal to 400 kilos): 200 - 300 mL via S.C.
- Cattle and Horses (200 - 300 kilos): 500 mL via E.V.
- Cattle (200 - 300 kilos): 50 to 80 mL via S.C.
- Sheep or Goats (25 - 60 kilos): 100 - 250 mL via E.V.
- Sheep or Goats (25 - 60 kilos): 40 mL via S.C. Pigs (greater than or equal to 120 kilos): 300 - 500 mL via E.V.
- Pigs (greater than or equal to 120 kilos): 300 - 500 mL via E.V. (Indicated in dehydration and bleeding).

## CONTRAINDICATIONS

---

Do not administer in pathological states that require low sodium intake.

## WARNINGS

---

Keep out of reach of children.

## GUARD PERIOD

---

**Meat:** 0 days.

**Milk:** 0 days.

## CONSERVATION

---

Store in a cool and dry place, at room temperature between 15° and 30°C., protected from sunlight.

## CONDITION OF SALE

---

To be supplied only on veterinary prescription.

## PRESENTATION

---

500 mL container

## PREPARED BY

---

Drag Pharma Laboratory.

## RECORDS

---

- Chile: Reg. SAG No. 1351
- Costa Rica: Reg. No. MAG CL4-19-01-4140
- Dominican republic: Reg. No. 5914
- Panamá: Reg. N° RF-4133-18

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian