

LOMBRIMIC® Oral suspension

BROAD SPECTRUM INTERNAL ANTIPARASITIC.



It is indicated for the treatment of mature and immature forms of gastrointestinal and pulmonary parasites and tapeworms, including ovicidal action.

Technical Specification

SPECIES

Bovine, sheep, goats, pigs and horses.

DOSAGE FORM

Oral suspension.

THERAPEUTIC ACTION

Broad spectrum internal antiparasitic.

COMPOSITION

Each 100 ml contains:

Fenbendazole.....10 g

Excipients q.s.p.....100 mL

INDICATIONS

It is indicated for the treatment of mature and immature forms of gastrointestinal and pulmonary parasites and tapeworms, including ovicidal action.

ACTION SPECTRUM

Bovine, sheep and goats: *Haemonchus spp.*, *Cooperia spp.*, *Ostertagia spp.*, *Trichostrongylus spp.*, *Oesophagostomum spp.*, *Strongyloides spp.*, *Trichuris spp.*, *Bunostomum spp.*, *Dictyocaulus spp.*, *Nematodirus spp.*, *Chabertia ovina*, *Moniezia spp.*

Horses: Large and small strongyles, *Parascaris equorum*, *Trichonema spp.*, *Trichostrongylus axei*, *Strongyloides*

westeri.

Pigs: *Oesophagostomum spp.*, *Metastrongylus spp.*, *Trichuris suis*, *Ascaris suum*, *Strongyloides ransomi*.

ROUTE OF ADMINISTRATION AND DOSAGE

Administer orally through a dosing gun or syringe.

Dose:

- **Cattle, sheep, goats, pigs:** 5 mL / 100 Kg of weight (Equivalent to 5 mg of Fenbendazole / Kg of weight). Single dose.
- **Horses:** 7.5 mL / 100 Kg of weight in a single dose (Equivalent to 7.5 mg of Fenbendazole / Kg of weight). Single dose.

CONTRAINDICATIONS

Do not administer in animals with known hypersensitivity to Fenbendazole.

SPECIAL PRECAUTIONS FOR THE OPERATOR

- Do not administer in animals with known hypersensitivity to Fenbendazole. Wear gloves when administering the product.
- Wash your hands after handling the product.
- Do not eat, drink or smoke while handling the product.
- Do not handle by people who are hypersensitive to Fenbendazole.
- Avoid contact with skin, eyes, or mouth.
- In case of accidental ingestion, go immediately to a medical center and show the product label.
- In case of contact with skin, eyes or mouth, wash immediately with plenty of water.
- In case of skin irritation, go immediately to a medical center and show the product label.

WARNINGS

Keep out of reach of children.

GUARD PERIOD

- **Cattle:** Meat: 14 days. Milk: 7 days
- **Horses:** Do not administer in animals whose meat is intended for human consumption
- **Sheep and goats:** Do not administer in animals whose milk is intended for human consumption.

OBSERVATIONS

Shake vigorously before use.

Special precautions for unused product or waste material:

- Empty containers can be disposed of as household waste, without any special precautions.
- Do not dispose of containers with product remains on the ground or water courses.
- For expired or unused products, contact the manufacturing laboratory.

CONSERVATION

Store between 15 and 30°C, protected from light.
Once the container is opened, use within 12 weeks. Discard unused product after that time frame.

CONDITION OF SALE

To be supply only on veterinary prescription.

PRESENTATION

Bottle with 100 ml, 500 ml, 1 L and 3 L

PREPARED BY

Drag Pharma Laboratory.

RECORDS

- Chile: Reg. SAG No. 0624
- Dominican Republic: Reg. No. 6516

You have entered drug info or veterinary products intended exclusively (s) kind (s) indicated (s).

Drag Pharma Lab is not responsible for the consequences of misuse of the products, and the use of this information without consulting a veterinarian